

## STA 711 Homework 5

**Due:** Friday, March 3, 12:00pm (noon) on Canvas.

**Instructions:** Submit your work as a single PDF. For this assignment, you may include written work by scanning it and incorporating it into the PDF. Include all R code needed to reproduce your results in your submission.

### Convergence of random variables

1. For each of the following sequences  $\{Y_n\}$ , show that  $Y_n \xrightarrow{p} 1$ .
  - (a)  $Y_n = 1 + nX_n$ , where  $X_n \sim \text{Bernoulli}(1/n)$
  - (b)  $Y_n = \frac{1}{n} \sum_{i=1}^n X_i^2$ , where  $X_i \stackrel{iid}{\sim} N(0, 1)$
2. Suppose that  $Y_1, Y_2, \dots \stackrel{iid}{\sim} \text{Beta}(1, \beta)$ . Find a value of  $\nu$  such that  $n^\nu(1 - Y_{(n)})$  converges in distribution.
3. Suppose that  $Y_1, Y_2, \dots \stackrel{iid}{\sim} \text{Exponential}(1)$ . Find a sequence  $a_n$  such that  $Y_{(n)} - a_n$  converges in distribution.
4. In this problem, we will prove part of the continuous mapping theorem. Let  $\{Y_n\}$  be a sequence of random variables such that  $Y_n \xrightarrow{p} Y$  for some random variable  $Y$ . Let  $g$  be a continuous function; recall that  $g$  is continuous if for all  $\varepsilon > 0$ , there exists some  $\delta > 0$  such that  $|g(x) - g(y)| < \varepsilon$  whenever  $|x - y| < \delta$ . Prove that  $g(Y_n) \xrightarrow{p} g(Y)$ .

### Normal distributions and the Wald test

Suppose that  $\hat{\theta}$  is some estimator of a parameter of interest  $\theta \in \mathbb{R}^d$ . We want to test the hypotheses

$$H_0 : \theta = \theta_0 \quad \text{vs.} \quad H_A : \theta \neq \theta_0.$$

If  $\hat{\theta}$  is approximately normal, then we can use the Wald test (often  $\hat{\theta}$  will be the MLE, but the Wald test can be applied to any asymptotically normal estimator, not just to the MLE). Formally, suppose that

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} N(0, V),$$

and let  $\hat{V}$  be some estimator of the covariance matrix  $V$ , such that  $\hat{V} \xrightarrow{p} V$ . Then the Wald test statistic is

$$W = n(\hat{\theta} - \theta_0)^T \hat{V}^{-1} (\hat{\theta} - \theta_0).$$

The goal of this section is to verify that  $W \xrightarrow{d} \chi_d^2$  if  $H_0$  is true. Our derivation will rely on the following properties of multivariate normal distributions, and positive semi-definite matrices:

- Recall from HW 4 that if  $X \sim N(\mu, \Sigma)$ , then

$$\mathbf{a} + \mathbf{B}X \sim N(\mathbf{a} + \mathbf{B}\mu, \mathbf{B}\Sigma\mathbf{B}^T)$$

- For any random vector  $X$ , the covariance matrix  $\Sigma = \text{Var}(X)$  is positive semi-definite (you may use this without proof)

- If  $\Sigma$  is a positive semi-definite matrix, then there exists a unique positive semi-definite matrix  $\Sigma^{\frac{1}{2}}$  such that  $\Sigma = \Sigma^{\frac{1}{2}} \Sigma^{\frac{1}{2}}$  (you may use this without proof)
- $Z \sim N(0, \mathbf{I})$  if and only if  $Z = (Z_1, \dots, Z_q)^T \stackrel{iid}{\sim} N(0, 1)$  (you may use this without proof).
- Suppose that  $X = (X_1, \dots, X_q)^T \sim N(\mu, \Sigma)$ . The entries  $X_i$  and  $X_j$  are independent *if and only* if  $\Sigma_{ij} = \text{Cov}(X_i, X_j) = 0$ . This is a special property of multivariate normal distributions, which we will prove below.
- If  $Z \sim N(0, \mathbf{I})$  is a  $q$ -dimensional multivariate normal variable, where  $\mathbf{I}$  is the identity matrix, then  $Z^T Z \sim \chi_q^2$  (we will prove this below).

5. Let us begin by proving some results for the multivariate normal.

- Show that if  $X \sim N(\mu, \Sigma)$ , then  $\Sigma^{-\frac{1}{2}}(X - \mu) \sim N(0, \mathbf{I})$ , where  $\mathbf{I}$  is the identity matrix.
- Show that  $X \sim N(\mu, \Sigma)$  if and only if  $X = \mu + \Sigma^{\frac{1}{2}} Z$  where  $Z \sim N(0, \mathbf{I})$ .
- Let  $X \sim N(\mu, \Sigma)$ , where  $X \in \mathbb{R}^q$ . Suppose that for some  $1 \leq p < q$ ,  $\Sigma$  can be partitioned as

$$\Sigma = \begin{pmatrix} \Sigma_{11} & 0_{p \times (q-p)} \\ 0_{(q-p) \times p} & \Sigma_{22} \end{pmatrix},$$

where  $\Sigma_{11}$  is  $p \times p$ ,  $\Sigma_{22}$  is  $(q-p) \times (q-p)$ , and  $0_{m \times n}$  denotes the matrix of zeros of the specified dimensions. Similarly partition

$$X = \begin{pmatrix} X_{(1)} \\ X_{(2)} \end{pmatrix} \quad \mu = \begin{pmatrix} \mu_{(1)} \\ \mu_{(2)} \end{pmatrix},$$

into vectors of length  $p$  and  $q-p$ . Prove that

$$X_{(1)} \sim N(\mu_{(1)}, \Sigma_{11}), \quad X_{(2)} \sim N(\mu_{(2)}, \Sigma_{22}),$$

and  $X_{(1)}$  and  $X_{(2)}$  are independent.

- Using (c), conclude that if  $X = (X_1, \dots, X_q)^T \sim N(\mu, \Sigma)$ , then the entries  $X_i$  and  $X_j$  are independent *if and only* if  $\Sigma_{ij} = \text{Cov}(X_i, X_j) = 0$ .

6. Now let's derive the relationship between the normal distribution and the  $\chi^2$  distribution.

- Let  $Z \sim N(0, 1)$  be a standard normal variable. Show that  $Z^2 \sim \chi_1^2$  (a  $\chi^2$  distribution with 1 degree of freedom), by proving that the pdf of  $Y = Z^2$  is

$$f_Y(y) = \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{y}} e^{-y/2}.$$

- Suppose that  $Z_1, Z_2, \dots, Z_q \stackrel{iid}{\sim} N(0, 1)$ . Show that  $\sum_{i=1}^q Z_i^2 \sim \chi_q^2$  (a  $\chi^2$  distribution with  $q$  degrees of freedom).
- Let  $\theta \in \mathbb{R}$  be a parameter of interest, and  $\hat{\theta}_n$  the maximum likelihood from a sample of size  $n$ . Let

$$Z_n = \sqrt{n\mathcal{I}_1(\theta)}(\hat{\theta}_n - \theta).$$

Asymptotic normality of the MLE tells us that  $Z_n \xrightarrow{d} N(0, 1)$ . Show that  $Z_n^2 \xrightarrow{d} \chi_1^2$ .

7. Finally, let's connect the multivariate normal with the  $\chi^2$ .

- (a) Show that if  $X \sim N(\mu, \Sigma)$ , then  $(X - \mu)^T \Sigma^{-1} (X - \mu) \sim \chi_q^2$ .
- (b) Suppose that  $\hat{\theta}$  is some estimator of  $\theta \in \mathbb{R}^d$ , and  $\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} N(0, V)$ . Let  $\hat{V}$  be an estimator of  $V$  such that  $\hat{V} \xrightarrow{p} V$ , and let  $W = n(\hat{\theta} - \theta_0)^T \hat{V}^{-1} (\hat{\theta} - \theta_0)$ . Prove that  $W \xrightarrow{d} \chi_d^2$  if the null hypothesis  $H_0 : \theta = \theta_0$  is true.