

# Probability review

This review sheet provides a summary of some of the important definitions and properties from probability which will be useful in STA 711. It is by no means complete. For full details, see Casella & Berger, chapters 1, 2, and 4.

## CDFs, density functions, and probability mass functions

- *Cumulative distribution function (cdf)*: Let  $X$  be a random variable. The cdf of  $X$  is defined by

$$F_X(x) = \mathbb{P}(X \leq x).$$

- $X$  is a *continuous* random variable if  $F_X(x)$  is a continuous function of  $x$ , and  $X$  is a *discrete* random variable if  $F_X(x)$  is a step function of  $x$ .
- *Probability mass function (pmf)*: The pmf of a discrete random variable  $X$  is  $f(x) = \mathbb{P}(X = x)$ .
- *Probability density function (pdf)*: The pdf of a continuous random variable  $X$  is the function which satisfies

$$F_X(x) = \int_{-\infty}^x f(x)dx.$$

## Joint, marginal, and conditional distributions

Let  $X$  and  $Y$  be two random variables.

- *Joint cdf*: The joint cdf of  $X$  and  $Y$  is

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y).$$

- *Joint mass function*: If  $X$  and  $Y$  are discrete, their joint mass function is  $f(x, y) = \mathbb{P}(X = x, Y = y)$ .
- *Joint pdf*: If  $X$  and  $Y$  are continuous, their joint pdf is the function  $f(x, y)$  such that for every set  $A \subset \mathbb{R} \times \mathbb{R}$ ,

$$\mathbb{P}((X, Y) \in A) = \int_A \int f(x, y) dx dy$$

- *Marginal distributions*: Given a joint pdf  $f(x, y)$ , the marginal pdf of  $X$  is given by

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$$

and the marginal pdf of  $Y$  is given by

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$$

(For discrete random variables, the definitions are similar, just replace integrals with sums)

- *Conditional distributions:* Given a joint pdf or pmf  $f(x, y)$ , the conditional pdf/pmf of  $X|Y = y$  is defined by

$$f(x|y) = \frac{f(x, y)}{f_Y(y)},$$

for any  $y$  such that  $f_Y(y) > 0$ .

## Probability, expectation, and variance

- *Expectation:* The *expectation*, or *mean*, of a random variable  $X$  is

$$\mathbb{E}[X] = \begin{cases} \sum x f(x) & X \text{ is discrete} \\ \int_{-\infty}^{\infty} x f(x) dx & X \text{ is continuous} \end{cases}$$

- *Variance:* The *variance*, or second central moment, of a random variable  $X$  is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}(X))^2]$$

- *Covariance:* If  $X$  and  $Y$  are two random variables, the *covariance* of  $X$  and  $Y$  is

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}(X))(Y - \mathbb{E}(Y))].$$

- *Conditional expectation:* The conditional expectation of  $X$  given  $Y = y$ , denoted  $\mathbb{E}[X|Y = y]$ , is given by

$$\mathbb{E}[X|Y = y] = \begin{cases} \sum x f(x|y) & X \text{ is discrete} \\ \int_{-\infty}^{\infty} x f(x|y) dx & X \text{ is continuous} \end{cases}$$

- *Law of total probability:* Let  $A$  be an event and  $B_1, \dots, B_k$  be disjoint event which partition the space (i.e,  $P(B_i \cap B_j) = 0$  if  $i \neq j$ , and  $\sum_i P(B_i) = 1$ ). Then,

$$P(A) = \sum_{i=1}^k P(A|B_i)P(B_i)$$

- *Law of total expectation* (aka law of iterated expectation):

$$\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X|Y]]$$

(Note here that  $\mathbb{E}[X|Y]$  is a random variable which is a function of  $Y$ ). We can apply this rule to conditional expectations, too:

$$\mathbb{E}[X|Y_1] = \mathbb{E}[\mathbb{E}[X|Y_1, Y_2]|Y_1]$$

- *Law of total variance* (aka law of iterated variance):

$$\text{Var}(X) = \mathbb{E}[\text{Var}(X|Y)] + \text{Var}(\mathbb{E}[X|Y])$$

## Functions of random variables

- *Law of the unconscious statistician:* Let  $X$  be a random variable with pdf or pmf  $f(x)$  (depending on whether  $X$  is continuous or discrete). Let  $g(X)$  be a function of  $X$ . Then

$$\mathbb{E}[g(X)] = \sum_x g(x)f(x) \quad X \text{ is discrete}$$

$$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx \quad X \text{ is continuous}$$

- *Finding the distribution of a transformation:* Let  $X$  be a continuous random variable with pdf  $f_X(x)$ , and let  $Y = g(X)$  be a function of  $X$ . To find the distribution of  $Y$ :

1. For each  $y$ , find the set  $A_y = \{x : g(x) \leq y\}$
2. Find the cdf:

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(g(X) \leq y) = \int_{A_y} f_X(x)dx$$

3. The pdf is  $f_Y(y) = \frac{d}{dy}F_Y(y)$

There is a special case when  $g$  is a monotone function. If  $X$  is continuous and  $g$  is monotone, then

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy}g^{-1}(y) \right|.$$

This special case can be extended if there exists a partition such that  $g$  is monotone on each piece of the partition (see Theorem 2.1.8 in Casella & Berger).

## Moment generating functions

- *Moments:* Let  $X$  be a random variable. The  $n$ th *moment* of  $X$  is  $\mathbb{E}[X^n]$ .
- *Moment generating function (mgf):* The mgf of  $X$  is  $M_X(t) = \mathbb{E}[e^{tX}]$ , provided that the expectation exists for  $t$  in some neighborhood of 0.
- *Properties of mgfs:*

$$(a) \text{ If } X \text{ has mgf } M_X(t), \text{ then } \mathbb{E}[X^n] = \frac{d^n}{dt^n} M_X(t) \Big|_{t=0}$$

- (b) If  $X$  and  $Y$  are independent, with mgfs  $M_X(t)$  and  $M_Y(t)$ , then the mgf of  $X + Y$  is

$$M_{X+Y}(t) = M_X(t)M_Y(t)$$

- (c) Let  $X$  and  $Y$  be random variables with cdfs  $F_X$  and  $F_Y$ . If  $M_X(t) = M_Y(t)$  for all  $t$  in an open interval around 0, then  $F_X(u) = F_Y(u)$  for all  $u$ .
- (d) Let  $a, b \in \mathbb{R}$ , and let  $Y = a + bX$ . The mgf of  $Y$  is

$$M_Y(t) = e^{at} M_X(bt).$$

## Statistics with matrix algebra

- *Definition of expectation and variance:* Let  $X = (X_1, \dots, X_k)^T \in \mathbb{R}^k$  be a random vector. Then

$$\mathbb{E}[X] = (\mathbb{E}[X_1], \dots, \mathbb{E}[X_k])^T,$$

and

$$\text{Var}(X) = \Sigma$$

where  $\Sigma \in \mathbb{R}^{k \times k}$  is the covariance matrix for  $X$ , with entries  $\Sigma_{ij} = \text{Cov}(X_i, X_j)$ . (This implies that the diagonal entries are  $\Sigma_{ii} = \text{Var}(X_i)$ ).

- *Expectation and variance of linear combinations:* Let  $X \in \mathbb{R}^k$  be a random vector, and let  $\mathbf{a} \in \mathbb{R}^k$  and  $\mathbf{B} \in \mathbb{R}^{m \times k}$ . Then

$$\mathbb{E}[\mathbf{a} + \mathbf{B}X] = \mathbf{a} + \mathbf{B}\mathbb{E}[X]$$

$$\text{Var}(\mathbf{a} + \mathbf{B}X) = \mathbf{B}\text{Var}(X)\mathbf{B}^T$$

- *Matrix square roots:* If  $M$  is a positive semi-definite matrix, then  $M^{\frac{1}{2}}$  is the unique positive semi-definite matrix such that  $M = M^{\frac{1}{2}}M^{\frac{1}{2}}$ . If  $M = \text{diag}(m_1, \dots, m_k)$ , then  $M^{\frac{1}{2}} = \text{diag}(\sqrt{m_1}, \dots, \sqrt{m_k})$ .
- *Block matrix inverses:* Let

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$$

be a block matrix with  $A \in \mathbb{R}^{p \times p}$ ,  $B \in \mathbb{R}^{p \times q}$ ,  $C \in \mathbb{R}^{q \times p}$ , and  $D \in \mathbb{R}^{q \times q}$ . Assuming that  $A$  and  $D$  are invertible, then

$$M^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1} \end{bmatrix}$$